arXiv:0803.0872v1 [cond-mat.stat-mech] 6 Mar 2008

Cluster formation in the Fermi system with long-range interaction

K.V. Grigorishin* and B.I. Lev†

Bogolyubov Institute for Theoretical Physics of the Ukrainian National Academy of Sciences,
14-b Metrolohichna str. Kiev-03680, Ukraine.

Based on statistical approach we described possible formation of spatially inhomogeneous distribution in the system of interacting Fermi particles by long-rage forces, and we demonstrated nonperturbative calculation of the partition function in this case. It was shown, that particles interacting with an attractive $1/r$ potential form clusters. Cluster is equilibrium structure, if we suppose that average energy of interaction of two particles is much less than their average kinetic energy kT. The analogy between self-gravitation gas and plasma was shown. The dynamics of cluster formation was considered with help hydrodynamical and statistical approaches, and time of relaxation to equilibrium state was found.

PACS numbers: 05.30-d, 05.70.Fh

I. INTRODUCTION.

The formation of spatially inhomogeneous distribution of interacting particles is a typical problem in condensed matter physics. The types of spatial structures and conditions for their formation are determined by type of interaction. A cluster is one from forms of spatial inhomogeneity. For gas with attraction of Coulomb type between particles (self-acting system) we can't calculate the virial coefficients (we cannot do it for interaction $1/r^n$ if $n \leq 3$ [27]). Statistical description of the such system developed in Refs. [1, 2, 3, 4, 5] is based on the application of the apparatus of quantum theory of field [12, 13, 14, 15, 16, 17, 23] where it was shown that gas with interacting particles is equivalent to two scalar fields corresponding attraction and repulsion accordingly with an exponential self-action. The partition function can be representation in terms of a functional integral over these auxiliary fields.

---

*Electronic address: konst@orta.net.ua

†Electronic address: blev@i.kiev.ua

The extremal conditions for this functional are nonlinear equations. The spatial distribution function, which describes the cluster, is soliton solution of the equations. This approach gives the possibility to find the spatial distribution of particles, to calculate the cluster's size and to determine the temperature of phase transition in to the state under consideration by nonperturbative calculation.

In the papers [6, 7], this approach has been applied to investigation of the system with long-range interacting Bose particles. It has been shown that spatial distribution of such system is inhomogeneous and has view of finite-size cluster. The radius of cluster, conditions and dynamics of its formation have been found both for undegenerated states and for Bose-Einstein condensation regime. In this article, based on statistical approach [1, 2, 3, 4], we demonstrate a nonperturbative calculation of the partition function, we solve the system of Fermi particles interacting with an attractive $1/r$ potential, that is we obtain free energy for the such system. We obtain expression for equilibrium radius of cluster and we described dynamics of the cluster formation. However, we can find exact results in Boltsman limit with first quantum corrections. For all temperatures we propose the approach which gives the possibility to find radius of the cluster approximately or to evaluate of one.

So far as the cluster's radius we can understand as size of spatial inhomogeneity of a system, then it is present interest to investigate the general properties of spatial distribution function of the systems with Coulomb interacting particles - above-mentioned gas of gravitating particles and the system of repulsing particles. This can be done in statistical and hydrodynamical approaches. Using hydrodynamical approaches we can find the times of relaxation of the systems with Coulomb interacting particles.

II. STATISTICAL APPROACH TO THE SYSTEM OF INTERACTING PARTICLES.

Let's consider an interacting particles' system being in such conditions when, on the one hand, wave's thermal length of a particle can be larger than average distance between them, so that it is necessary to take into account type of the statistic, but, on the other hand, this length is by far smaller than average scattering length, that lets to describe interaction classically disregarding dynamical quantum correlations. The Hamiltonian of such a system [1, 3, 17, 18] is

$$H(n) = \sum_s \varepsilon_s n_s - \frac{1}{2}\sum_{ss'} W_{ss'} n_s n_{s'} + \frac{1}{2}\sum_{ss'} U_{ss'} n_s n_{s'}\,, \tag{1}$$

where ε_s is the additive part of particle energy in the state s (for example kinetic energy or energy in an external field), $W_{ss'}$ and $U_{ss'}$ are the absolute values of the attraction and repulsion energies of particles in the states s and s' respectively. The macroscopic state of the system is determined by occupations numbers n_s. The subscript s corresponds to variables that describe an individual particle state.

In Refs.[1, 2, 3, 4] in order to investigate thermodynamical properties of the interacting particles' system a Hubbard-Stratonovich [8, 9, 10, 11] representation has been used for the partition function in the terms of grand partition function

$$Z_n = \frac{1}{2\pi i}\oint d\xi \int D\varphi \int D\psi \exp\left(-S\left(\xi,\varphi,\psi\right)\right)\,, \tag{2}$$

where S is the functional which we call an effective free energy which is analogous to an action in field theory:

$$\begin{aligned} S\left(\xi,\varphi,\psi\right) = \quad & \frac{1}{2\beta}\sum_{s,\acute{s}}\left(W^{-1}_{s,s'}\varphi_s\varphi_{s'} + U^{-1}_{s,s'}\psi_s\psi_{s'}\right) \\ & +\delta\sum_s \ln\left(1 - \delta\xi\exp\left(-\beta\varepsilon_s + \varphi_s\right)\cos\psi_s\right) + (N+1)\ln\xi\,, \end{aligned} \tag{3}$$

$\xi \equiv e^{\beta\mu}$ is activity, μ is chemical potential, $\beta = 1/kT$ is reverse temperature, s and $s^{'}$ run all the states of the system, ε is kinetic energy, N is number of particles, $\delta = +1$ for Bose particles and -1 for Fermi particles. The two auxiliary fields φ and ψ are introduced corresponding to attraction and repulsion. The partition function (2) is written as a functional integral over these fields. $W^{-1}_{s,s^{'}}, U^{-1}_{s,s^{'}}$ are inverse operators of the interaction: $\omega^{-1}_{ss^{'}} = \delta_{ss'}\hat{L}_{s'}$ where $\hat{L}_{s'}$ is such an operator for which the interaction's potential is Green function.

The integral (2) is calculated by the method of "saddle-point" [19, 20] across the point determined by the functional derivatives $\frac{\delta S}{\delta\varphi} = \frac{\delta S}{\delta\psi} = 0$ as

$$\frac{1}{\beta}\sum_{s'} W^{-1}_{ss'}\varphi_{s'} - \frac{\xi_s e^{\varphi_s}\cos\psi_s}{1 - \delta\xi_s e^{\varphi_s}\cos\psi_s} = 0 \tag{4}$$

$$\frac{1}{\beta}\sum_{s'} U^{-1}_{ss'}\psi_{s'} + \frac{\xi_s e^{\varphi_s}\sin\psi_s}{1 - \delta\xi_s e^{\varphi_s}\cos\psi_s} = 0 \tag{5}$$

and the derivative

$$\frac{\partial S}{\partial \xi} = 0 \Rightarrow \sum_s \frac{\xi_s e^{\varphi_s}\cos\psi_s}{1 - \delta\xi_s e^{\varphi_s}\cos\psi_s} = N+1\,, \tag{6}$$

where $\xi_s = \xi e^{\beta\varepsilon_s}$. This set of equations (4-6) provides a solution of the many-particle problem in the sense that it selects the system states whose contributions in the partition function are dominant. The third equation is normalization condition for the particle distribution function which is determined by the auxiliary fields. It is obvious that, for given statistics, the distribution function depends on the interaction nature and intensity. The cluster formation corresponds to particle localization within a limited space. In our treatment, the effect is reflected in the behavior of the auxiliary fields and chemical potential.

For the screened Coulomb or Newtonian potential, the inverse operator may be written as

$$\widehat{L} = -\frac{1}{4\pi q^2}(\triangle - \lambda^2), \tag{7}$$

where q^2 is the interaction constant, $\triangle$ is Laplace's operator and λ^{-1} is the screening length [8, 15, 16, 17]. The number of realistic interactions, for which the inverse operator can be found and general solution of the set (4-6) can be obtained, is limited. That's why we'll confine ourselves to long-range both attractive and repulsive Coulomb potential ($\propto \frac{1}{R}$) for Fermi gas where we'll find the conditions for the cluster formation and cluster parameters.

In the continuum approximation, the subscript s runs through a continuum of values in the system of volume V. When integrating over impulses and coordinates, we bear in mind that the unit of cell's volume in the space of individual states is equal to $\omega = (2\pi\hbar)^3$. In order to avoid unnecessary complications we shall consider particles without spin only.

III. CLUSTER FORMATION IN THE SYSTEM OF ATTRACTING PARTICLES.

A. The equations for spatial distribution function for system with Coulomb attraction.

Now we shall consider the system of particles interacting by long-range attraction only and we shall demonstrate a nonperturbative calculation of the partition function. For Coulomb attraction the inverse operator is known to be $W^{-1}_{rr'} = \frac{-1}{4\pi q^2}\triangle_r \delta_{rr'}$, where g is the interactional constant and $\triangle_r$ is the Laplace operator.

Let's consider the system in the usual thermodynamical limit: number of particles $N \rightarrow \infty$ and volume $V \rightarrow \infty$ with fixed N/V. We consider the effective free energy (3) in spherical coordinates: $\nabla \equiv \frac{\partial}{\partial R}$ and, neglecting surface contribution, we can write the effective free energy (3) for Fermi gas as

$$\begin{aligned} S = \ & \frac{1}{2\beta}\int_0^V \frac{(\nabla\varphi)^2}{4\pi q^2}dV - \frac{1}{\omega}\int_0^V dV\int d^3p\ln\left[1+\xi e^{\varphi}\exp\left(-\beta\frac{p^2}{2m}\right)\right] \\ & +N\ln\xi = \int_0^V dV\left[\frac{(\nabla\varphi)^2}{4r_q} - \frac{1}{\lambda^3}f_{5/2}\left(\xi e^{\varphi}\right)\right] + N\ln\xi, \end{aligned} \tag{8}$$

where $r_q = 2\pi q^2\beta$, m is a particle mass, $\lambda = \sqrt{\frac{\beta\hbar^2}{2\pi m}}$ is wave's thermal length of a particle and $f_{5/2}\left(\xi\right)$
$= \frac{4}{\sqrt{\pi}}\int_0^\infty dx x^2\ln\left(1+\xi e^{-x^2}\right) = \sum_{l=1}^{\infty}(-1)^{l+1}\frac{\xi^l}{l^{5/2}}$ is special Fermi function [19, 20]. Sense of the auxiliary field φ is next - a spatial distribution's function can be expressed by φ as

$$\rho(R) = \int_0^\infty \frac{d^3p}{(2\pi\hbar)^3}\frac{\xi\exp\left(-\frac{p^2}{2mkT}\right)e^{\varphi(R)}}{1+\xi\exp\left(-\frac{p^2}{2mkT}\right)e^{\varphi(R)}}, \tag{9}$$

where we used the underintegral expression in Eq.(6). Let's introduce the dimensionless quantity $r = R/r_q$ instead of R. Then, the effective free energy (8) (in spherical coordinates) can be written as

$$S = 4\pi\int_0^V r^2dr\left[\frac{(\nabla\varphi)^2}{4} - \frac{r_q^3}{\lambda^3}f_{5/2}\left(\xi e^{\varphi}\right)\right] + N\ln\xi. \tag{10}$$

The saddle-point equation is Lagrange equation for this functional:

$$\frac{\partial^2\varphi}{\partial r^2} + \frac{2}{r}\frac{\partial\varphi}{\partial r} + \frac{2r_q^3}{\lambda^3}\frac{\partial f_{5/2}\left(\xi e^{\varphi}\right)}{\partial\varphi} = 0. \tag{11}$$

This equation selects the system's states (which are described by the field configuration $\varphi(R)$ or spatial distribution function $\rho[\varphi(R)]$) whose contributions in the partition function are dominant. Unfortunately, this equation hasn't analytical solution. But the problem simplifies in Boltzmann limit.

B. The solution in Boltzmann limit and dynamics of the cluster formation.

Let's consider the limit case $\xi \to 0$ corresponding to high temperature and small concentration, using decomposition of the special function $f_{5/2}\left(\xi e^{\varphi}\right)$ in row on orders of the activity ξ. Then the effective free energy(8) is reduced to the simpler expression

$$S = 4\pi\int_0^V\left[\frac{1}{4}(\nabla\varphi)^2 - \xi\frac{r_q^3}{\lambda^3}e^{\varphi}\right]r^2dr + N\ln\xi. \tag{12}$$

That is the system with Coulomb attraction is equivalent to single scalar field $\varphi(r)$ with an exponential self-action. Analogous expression was obtained in Ref.[23], however therm $N\ln\xi$, which fixes number of particles, is absent there. The saddle-point equation for the functional (12) is

$$\frac{\partial^2\varphi}{\partial r^2}+\xi\frac{2r_q^3}{\lambda^3}e^{\varphi}=0\,. \tag{13}$$

As it will be shown below, the term $\frac{2}{r}\frac{\partial\varphi}{\partial r}$ can be omitted. In order to connect the auxiliary field φ with the density $\rho(R)$ we have to use Eq.(9) and to pass to the Bolzmann limit $\xi\longrightarrow 0$:

$$\rho=\frac{\xi}{\lambda^3}e^{\varphi}\equiv\frac{\xi}{\lambda^3}\sigma^2, \tag{14}$$

where we introduce the new variable $\sigma=\exp\left(\varphi/2\right)$ and let's mark in $\alpha^2\equiv r_q^3/\lambda^3$. Then we can rewrite Eq.(13) for the field φ as equation for the density as

$$\frac{\partial^2\sigma}{\partial r^2}-\frac{1}{\sigma}\left(\frac{\partial\sigma}{\partial r}\right)^2+\xi\alpha^2\sigma^3=0\,. \tag{15}$$

This equation has a soliton solution [3]

$$\sigma=\frac{\Delta}{\sqrt{\xi}\alpha}\frac{1}{\cosh\Delta r}, \tag{16}$$

where Δ is an integration constant. Any soliton solution corresponds to a spatially inhomogeneous distribution of particles - a finite-size cluster. The corresponding asymptotics are $\sigma^2=1$ for $r=d$, where d is the cluster size, and $\sigma\longrightarrow 0$ as $r\longrightarrow\infty$. This solution describes the presence of particles in the inhomogeneous formation of the size d and the absence of particles at infinity (Fig.1). This spatial distribution is compressing to line $\sigma=1$ or $\varphi=0$ when $T\longrightarrow\infty$, $\frac{N}{V}\longrightarrow 0$, that's why the field $\varphi=0$ corresponds to spatially homogeneous distribution in the system.

Let's substitute solution (16) in the effective free energy (12):

$$S=4\pi\int_0^d\left(\Delta^2-2\xi\alpha^2\sigma^2\right)r^2dr+N\ln\xi\,. \tag{17}$$

Then we'll integrate using the decomposition $1/\cosh x\approx 1-x^2/2$ in power series of $x\equiv\Delta d<<1$:

$$S=-V\frac{\Delta^2}{\alpha^2\lambda^3}+N\ln\xi\,. \tag{18}$$

The integration constant Δ^2 is being found from the asymptotics $\sigma^2(d)=1$: $\Delta^2\approx\xi\alpha^2+\xi^2d^2\alpha^4$. Thus, we have the result:

$$S=-\frac{V}{\lambda^3}\xi+N\ln\xi-\frac{V}{\lambda^3}\xi^2d^2\alpha^2\,. \tag{19}$$



FIG. 1: The spatial distribution function $\sigma(r)^2$ in a cluster at different temperatures (schematically). The solid line corresponds to a lower temperature T_1, the dash line corresponds to a higher temperature T_2. The dot line $\sigma^2 = 1$ corresponds to mean fixed density of the system and determines equilibrium radii of cluster d_{01} and d_{02} under the above-mentioned thermodynamical conditions.

Assuming that the average energy of interaction of two particles is less than the average kinetic energy $\sim kT$ of a particle $\frac{r_q^3 N}{V} \ll 1$, we can find the activity ξ from the equation $\frac{\partial S}{\partial \xi} = 0$ as

$$\xi \approx \frac{\lambda^3 N}{V} - \frac{2d^2\alpha^2\lambda^6 N^2}{V^2} \equiv \xi_0 + \xi_q \,, \;\; |\xi_q| \ll \xi_0 \,, \tag{20}$$

where ξ_0 and ξ_q are activities of ideal gas and first correction for the interaction accordingly. Then integrating Eq.(2) with the effective free energy (19) on saddle-point (20) we obtain the partition function as

$$Z_N = Z_N^0 \times \exp\left[\frac{V}{\lambda^3}\xi_q - N\ln\left(1 + \frac{\xi_q}{\xi_0}\right) + \frac{V}{\lambda^3}\xi^2 d^2\alpha^2\right] , \tag{21}$$

where Z_N^0 is the partition function for ideal gas. Knowing it, we can find a decomposition of free energy on cluster's radius d which plays a part correlation length

$$\begin{aligned} F &= F_0 - kT\left[\frac{V}{\lambda^3}\xi_q - N\ln\left(1 + \frac{\xi_q}{\xi_0}\right) + \frac{V}{\lambda^3}\xi^2 d^2\alpha^2\right] \\ &= F_0 + kT\left[-\frac{\alpha^2\lambda^3 N^2}{V}d^2 + \frac{\alpha^4\lambda^6 N^3}{V^2}d^4\right] , \end{aligned} \tag{22}$$

where F_0 is free energy for ideal gas. We can see that the d^2 term is always negative $-\frac{\alpha^2\lambda^3 N^2}{V} < 0$, that is cluster exists in the system with Coulomb attraction at any thermodynamical conditions. Minimizing (22) by the size of the cluster

$$\frac{\partial F}{\partial d} = -kT\frac{d\alpha^2\lambda^3 2N^2}{V}\left[1 - 4\frac{d^2\alpha^2\lambda^3 N}{V}\right] = 0\,, \tag{23}$$

we obtain the optimum radius of the cluster,

$$d_0^2 = \frac{V}{4Nr_q^3} \tag{24}$$

or, for the dimension value $D = d \cdot r_q$,

$$D_0^2 = \frac{1}{4}\frac{kT}{2\pi q^2}\frac{V}{N}\,. \tag{25}$$

This expression means that equilibrium size of the cluster is defined by balance of the two forces. First of them is attraction Coulomb energy, which aspires to compress the gas. It is represented by the multiplier q^2 (in sense of interaction constant). The decrease of the cluster's size with the increase of average density in the system N/V is connected with a closer packing of the particles in the cluster on account of the increase of the energy of interaction. The second of them is the thermal energy, which creates positive pressure resisting to the compression. It is represented by the multiplier kT.With increase of temperature the size of the cluster d is rising, however the central density is decreasing.When $T \to \infty$ the radius of the cluster $d \to \infty$ and the central density $\sigma(0) \to 1$. Such behavior the value d tells us that d plays a part of correlation length which in the point of phase transition (in our case is $T \to \infty$) in infinitely large. Such situation is realized due to the long-range type ($\sim 1/r$) of Coulomb interaction.

On other hand Eq.(25) may be understood as such such distance in the system on which the essential deflection from the average fixed density N/V is observed. If we assuming that $q^2 = Gm^2$, where m is particle mass and G is gravitational constant, this expression may be understood as Jeans length [22].

Let's consider the dynamics of cluster's formation. For it we'll use the equation of motion as

$$\frac{\partial d}{\partial t} = -\chi\frac{\partial F}{\partial d}, \tag{26}$$

where χ is a some constant. Applying (22) and (24) we have:

$$\dot{d} + \eta d^3 - \eta d_0^2 d = 0, \tag{27}$$

where we marked in $\eta \equiv \frac{\chi N k T}{2 d_0^4}$. The solution of Eq.(27) is

$$d^2 = \frac{d_0^2}{1 + C \exp\left(-2\eta d_0^2 t\right)}, \tag{28}$$

where C is integration constant. Let's consider the next initial states.

- Let the radius of a spatial inhomogeneity was larger than the equilibrium size $d(t) > d_0$ in some moment of time $t > 0$. Then the constant $-1 < C < 0$, that means the cluster's increase from spatially homogeneous distribution (where $d \to \infty$) to the equilibrium size d_0.

- Let the radius of a spatial inhomogeneity was smaller than the equilibrium size $d(t) < d_0$. Then the constant $C > 0$, that that means the cluster's evaporation from initial distribution with central density which is larger then equilibrium density at the given temperature.

We can see that size of the cluster approaches to the equilibrium size (24) asymptotically $d = d_0[1 + C \exp(-2\eta d_0^2 t)]$ when deflections are small. The term $1/2\eta d_0^2$ can be understood as time of relaxation. If some fluctuation of density has appeared in the system, then it brings about the appearance of the potential's gradient. In its turn, it brings about the spatial inhomogeneity - cluster with the size approaching to the equilibrium value asymptotically. In others words, cluster is equilibrium structures if we are assuming that the average energy of interaction of two particles is less than the average kinetic energy $\sim kT$ of a particle $\frac{r_q^3 N}{V} \ll 1$, when the term $\frac{2}{r}\frac{\partial \varphi}{\partial r}$ in Eq.(13) can be omitted, as it will be in the next subsection.

Since the number of particles in the system $N \to \infty$, but the number of the particles in the cluster is finite, then this means that our system disintegrates to the infinity multitude of clusters of the size D_0 each. After that the process of cluster formation repeats again, where the early formed clusters plays a part of particles. In others words, the free anergy of such system hasn't absolute minimum and each state of the system is analogous to false vacuum in theory of field [12].

1. *Single-particle motion.*

Let's consider single-particle motion in order to ground our assumptions in the previous section for calculation of cluster's size. For evaluation of cluster's size we can make use the another method without calculation of the free energy. Let's we assume that the average energy of Coulomb interaction of two particles is less than the average kinetic energy $\sim kT$ of a particle, when $\frac{r_q^3 N}{V} \ll 1$ that has been named by "local ideality approach" [24]. This must lead to the two consequences:

- the activity must be as
$$\xi = \xi_0 \left(1 + O\left[\frac{r_q^3}{V/N}\right]\right), \tag{29}$$
where $\xi_0 = \frac{\lambda^3 N}{V}$ is activity for ideal gas. That is the system's activity differences from activity for ideal gas little.

- The center density (Fig.1) differences from the average density little: $\rho(0) \simeq mN/V$. Then we can think that $\Delta^2 \simeq \frac{N}{V} r_q^3$ and we can evaluate the term $\frac{2}{r}\frac{\partial \varphi}{\partial r}$ or $\frac{2}{r}\frac{\partial \sigma}{\partial r}$ in Eqs.(13,15) with help the solution (16):
$\frac{\max\left|\frac{2}{r}\frac{\partial \sigma}{\partial r}\right|}{\frac{\partial^2 \sigma}{\partial r^2} - \frac{1}{\sigma}\frac{\partial \sigma}{\partial r} + \xi \frac{r_q^3}{\lambda^3}\sigma^3} \sim \frac{r_q^3 N}{V} \ll 1$. That is this term can be omitted in the local ideality approach.

After that let's calculate potential energy of a particle $U(r)$ in the cluster using Boltzmann distribution
$$\sigma(r) = \sigma(0) \exp\left[-\frac{U(r)}{kT}\right]. \tag{30}$$
Then using (16) we have that
$$U(r) = \quad -kT \ln\left[\frac{\sigma(r)}{\sigma(0)}\right] = 2kT \ln\left[\cosh\left(\Delta r\right)\right]. \tag{31}$$
The field $U(r)$ is self-coordinated field of the rest particles in the system (in the cluster). Using the assumption (29) about interactional and thermal energies and considering small distances from the center of the cluster as $r/r_q \leq 1$, we can write that
$$U(r) \approx kT\Delta^2 r^2 \approx \frac{kTNr_q^3}{V} r^2. \tag{32}$$
We can see that a particle in the cluster is in potential well created by all cluster's particles. Then cluster's radius d can be understood as such distant from center of the cluster that
$$U(d) = \frac{1}{2}kT, \tag{33}$$



FIG. 2: The potential energy $U(r)$ of a particle in the cluster (schematically). The solid line corresponds to a lower temperature T_1, the dash line corresponds to a higher temperature T_2. The particle does finite motion in the potential well created by the all particles in the system. Then the radius of the cluster d we determine as distance from the cluster's center to the point of stop of a particle with average kinetic energy $\frac{1}{2}kT$.

where, the multiplier 1/2 corresponds to radial degree of freedom. This means that d is distance from the cluster's center to the point of stop of a particle with average kinetic energy $\frac{1}{2}kT$ (Fig.2). That is radius of the cluster we calculate by self-coordinated manner.

Then solving the equation (33) we has the equilibrium radius of the cluster as $d^2 = \frac{V}{2r_q^3 N}$, which coincides with Eq.(24) with exactness to constant multiplier. Since two particles attracting by low $1/r$, they can not fall on each other in Newton mechanics. Hence spatial distribution is regular everywhere in the limits of the local ideality approach, however it is not so in the total case [25]. Such mechanism of cluster formation we shall call by the thermal mechanism. It is necessary to notice that single-particle motions in cluster is superdiffusive [26].

C. The cluster's size at all temperatures.

In order to calculate radius of a cluster at any temperatures we must solve Eq.(11) and obtain spatial distribution function of type (16). We can't do this by cause mathematical difficulties, however, if we suppose that gas is ideal locally, that is $\frac{r_q^3 N}{V} \ll 1$, and considered small distances from the coordinates' center $R/r_q \leq 1$ where $|\varphi| \ll 1$ (see Eq.(14) and Eq.(16)), then our problem simplifies.

For example, let's consider Eq.(13) and decompose e^{φ} in a row:

$$\frac{\partial^2\varphi}{\partial r^2} + \xi_0 \frac{2r_q^3}{\lambda^3} + O(\varphi) = 0, \tag{34}$$

where we suppose that $\xi \approx \xi_0$. Its solution and corresponding density is

$$\varphi \approx -\xi_0 \frac{2r_q^3}{\lambda^3} r^2 \Rightarrow \rho = \frac{\xi}{\lambda^3} \exp\left(-\frac{\xi_0 r_q^3}{\lambda^3} r^2\right). \tag{35}$$

Then we can obtain potential energy of a particle in the cluster as

$$U(r) = -kT \ln\left[\exp\left(-\frac{\xi_0 r_q^3}{\lambda^3} r^2\right)\right] = kT \frac{\xi_0 r_q^3}{\lambda^3} r^2, \tag{36}$$

which has be obtained early (32). Then, using the condition $U(d) = \frac{1}{2}kT$ and expression for the Boltzmann activity $\xi_0 = \frac{\lambda^3 N}{V}$ we obtain result

$$kT \frac{\xi_0 r_q^3}{\lambda^3} d^2 = \frac{1}{2}kT \Rightarrow d^2 = \frac{V}{2r_q^3 N}, \tag{37}$$

which coincides with Eq.(24).

Now let's to calculate the radius of cluster for any temperatures using Eq.(11) and decomposing one in a row as

$$\frac{\partial^2\varphi}{\partial r^2} + \frac{2r_q^3}{\lambda^3} \left(\frac{\partial f_{5/2}\left(\xi_0 e^{\varphi}\right)}{\partial \varphi}\right)_{\varphi=0} + O(\varphi) = 0. \tag{38}$$

Analogously, the size of the spatial inhomogeneity is determined by the term at zero degree of the field φ. However, the calculations with spatial distribution function (9) is somewhat problematical because the underintegral expression can not be integrated in elementary function. That's why we have to find another method. If we suppose that gas is ideal locally, that we have to suppose that motion of particles is quasiclassical [21]. Than we propose to use the particle's energy in Boltzmann view

$$U(r) = -kT \ln\left[\frac{\rho(r)}{\rho(0)}\right] = -kT \ln e^{\varphi(r)} = -kT\varphi(r), \tag{39}$$

where the distribution (14) has been used and the field φ is solution of the Eq.(38), however the radius of the cluster d we determine as

$$U(d) = \frac{1}{2}kT + \frac{3}{5}\varepsilon_F, \tag{40}$$

where ε_F is Fermi energy which is determined by the fixed average density of the system (A7). The term $\frac{1}{2}kT$ is a particle's kinetic energy caused by thermal motion and $\frac{3}{5}\varepsilon_F$ is average particle's kinetic energy in degenerated Fermi gas (because all particles can not be in the level $\varepsilon = 0$ by force of Pauli principle). The term $\frac{3}{5}\varepsilon_F$ can be understood as repulsive "statistical potential" [19] too. Then the assumption about local ideality of gas is written as $\frac{q^2}{\sqrt[3]{V/N}} \ll kT + \varepsilon_F$ - the average energy of interaction of two particles is less than the average kinetic energy of a particle.

Using the decomposition (38), equation (40) and equation (A3) for determination of the activity ξ_0 we can obtain expression for radius of the cluster for Coulomb interacting fermi-gas:

$$\left[\begin{array}{c} kT\frac{2r_q^3}{\lambda^3}\left(\frac{\partial f_{5/2}(\xi_0 e^{\varphi})}{\partial\varphi}\right)_{\varphi=0} d^2 = \frac{1}{2}kT + \frac{3}{5}\varepsilon_F \\ \\ \frac{\lambda^3 N}{V} = f_{3/2}(\xi_0) \end{array}\right]. \tag{41}$$

Let's consider some limit cases. In the limit of high temperatures we supposed that $kT \gg \varepsilon_F$. Then calculation the first quantum correction only in the decomposition (A3) we can obtain that

$$d^2 = \frac{V}{2r_q^3 N} + \frac{1}{2^{5/2}}\frac{\lambda^3}{r_q^3}, \tag{42}$$

where $\lambda/r_q \ll 1$. For the dimension value:

$$D^2 = \frac{kTV}{4\pi q^2 N} + \frac{1}{2^{7/2}}\frac{\hbar^3}{\pi(2\pi m)^{3/2}(kT)^{3/2}}. \tag{43}$$

The sense of the first addendum has been explained in the section (III B). The sense of the first quantum correction is explained below.

It is know that so-called "statistical potential" of particles' interaction exists in quantum gases [19]. It is repulsion for fermions and it is attraction for bosons. This phenomenon is connected with symmetry of particles' wave function. Hence the equilibrium size is determined by three energies - of Coulomb interaction (the interaction constant is r_q or q^2), thermal kT and above-mentioned "statistical potential" (λ or $\hbar$ plays a part of interaction constant). That is, in case of Fermi statistics, the quantum energy resists to the compression under action of Coulomb force as the small addendum to thermal energy for high temperatures. This explains the sign "+"for the small correction to the cluster's radius. For case Bose statistic we can analogous expression but the sign "+" replaced by the sign "−".

In the limit of low temperatures $kT \ll \varepsilon_F$ we obtain that

$$D^2 = 0.4\frac{\hbar^2}{q^2 m}\left(\frac{V}{N}\right)^{1/3}\left[1 + \frac{5}{6}\frac{kT}{\varepsilon_F}\right]. \tag{44}$$

We can see that the equilibrium size is determined by two energies - Coulomb potential and the repulsion statistical potential (the constant of interaction is $\hbar$). That is, the aforesaid quantum energy resists to the Colombian compression. The small correction $\frac{kT}{\varepsilon_F}$ means that above-mentioned thermal mechanism of cluster formation (as small addendum to Fermi statistical repulsion) including if $T \neq 0$.

Expression for radius of a cluster at low temperatures can be obtained by the next formal means. It is not difficult to notice, that the equilibrium radius of a cluster is determined by the multiplier $\frac{2r_m}{\lambda^3}$ in the equation (13) after the term e^{φ} at $\xi \approx \xi_0$. In the case $T \to 0$ we can suppose that the radius is determined by above-mentioned multiplier, however because temperature is zero, hence kT can not enter into the searching formula. Then it is necessary to multiply this coefficient to the value $\frac{kT}{\varepsilon_F}$ which cancellates temperature. Then the cluster's radius is determined as $\frac{1}{D^2} \propto \frac{r_q}{\lambda^3}\left(\frac{kT}{\varepsilon_F}\right) = \frac{\hbar^2}{q^2 m}\left(\frac{V}{N}\right)^{1/3}$, and we have the formula (44).

IV. FORMATION OF STRUCTURES IN INTERACTING PARTICLE'S SYSTEM WITH COULOMB REPULSION

A. Statistical approach

Let N_+ particles repulse by Coulomb low with the interaction constant q^2. In order to ensure stability of such system, we suppose that these particles is dipped in environment of contrary sign (with charge of $-q$), so that the condition of neutralization is executed: $N_+ = N_- = N$, but a current created by N_+ particles only. In other worlds, we consider such plasma where constant of interaction is q^2 and charge's carriers of one from the signs is immovable. Let's find spatial distribution function for such system, where $N \to \infty, V \to \infty$ and N/V is fixed.

Let a particle with charge q is situated in the beginning of coordinates. It polarizes the gas of particle with the same charge (the environment with contrary sign is immovable). For repulsion particles the action (3) has view:

$$S = \int_0^V \left[\frac{(\nabla\psi)^2}{4r_q} - \frac{1}{\lambda^3} f_{5/2}(\xi\cos\psi)\right] dV + N\ln\xi, \tag{45}$$



FIG. 3: The density $\rho(R)$ of polaron at different temperatures (schematically). The solid line corresponds to a lower temperature (T_1), the dash line corresponds to a higher temperature (T_2). The dot line $\rho = N/V$ corresponds to average fixed density.

and corresponding Lagrange equation is

$$\frac{d^2\psi}{dR^2} + \frac{2r_q}{\lambda^3}\frac{\partial f_{5/2}\left(\xi\cos\psi\right)}{\partial\psi} = 0. \tag{46}$$

The term $\frac{2}{r}\frac{\partial\psi}{\partial r}$ can be omitted as in the case of attracting particles. In Boltzmann limit $\xi \to 0$ we can write the action and spatial distribution function as

$$S = \int_0^V \left[\frac{(\nabla\psi)^2}{4r_q} - \frac{\xi}{\lambda^3}\cos\psi\right] dV + N\ln\xi, \tag{47}$$

$$\rho(R) = \frac{\xi}{\lambda^3}\cos\psi, \tag{48}$$

with corresponding Lagrange equation

$$\frac{d^2\psi}{dR^2} - \frac{2r_q}{\lambda^3}\xi\sin\psi = 0. \tag{49}$$

We assume that average energy of interaction of two particles is much less than their average kinetic energy, that is $\xi \approx \xi_0 = \frac{\lambda^3 N}{V}$, and we find solution in the area $\psi(R) \in [0, \pi/2]$ where $R \in [0, +\infty]$ with asymptotics: $\psi(0) = \pi/2$ and $\psi(+\infty) = 0$. The first of the asymptotics corresponds to absence of particles in the beginning of coordinates, because in order to penetrate there a particle must has infinite kinetic energy. Second of them means that the concentration equals to average concentration on large distances from the center $\rho(R \to \infty) = N/V$. In other words, the spatial distribution function must be pit - such formation called as polaron usually, unlike the system with attraction where spatial distribution is hump - cluster. In other words polaron can be called by anticluster. Corresponding solution of Eq.(49) is

$$\tan\frac{\psi}{4}=\tan\frac{\pi}{8}\exp\left(-\sqrt{\frac{2\xi_0 r_q}{\lambda^3}}R\right). \tag{50}$$

Then spatial distribution function (Fig.3) is written as

$$\rho(R)=\frac{N}{V}\cos\left[4\arctan\left(\exp\left(-\sqrt{\frac{2\xi_0 r_q}{\lambda^3}}R\right)\tan\frac{\pi}{8}\right)\right]. \tag{51}$$

This distribution is determined by balance of two energies: Coulomb energy which prevents from penetration of a particle in the center of coordinates, and the thermal energy (average kinetic particle's energy). Then radius of the polaron we can find as distance from its center to the point of stop: $U(D_p)\sim kT$. The function $U(r)$ we can find from Boltzmann distribution

$$\rho(R)=\rho(R\to\infty)\exp\left(-\frac{U(R)}{kT}\right). \tag{52}$$

It is using (50) we have

$$U=-kT\ln\left[\cos\left[4\arctan\left(\exp\left(-\sqrt{2\xi_0\frac{r_q}{\lambda^3}}R\right)\tan\frac{\pi}{8}\right)\right]\right]. \tag{53}$$

In order to find radius of the polaron let's apply the equation $U(D_p)=\frac{1}{2}kT$ and we obtain that

$$D_p^2\propto\frac{kTV}{4\pi q^2 N}. \tag{54}$$

This expression for polaron's radius coincides with cluster's radius (25) with exactness to the constant multiplier. This value we can understand as the size on which the neutralization of gas is violated.

The coincidences sizes D and D_p we can explain by the next way. We have seen that the spatial distributions in the both cases is result of balance of two energies only: Coulomb type and thermal energy. The radii of cluster and polaron are determined by the points of stop of a particle in the field of the rest particles. That is mechanism of formation of such structures is one and the same - a balance of they two energies. They are differed by roles only: in the gas of attracting particles they make spatial distribution of view "hump" and in the case of repulsing particles - "pit" Fig.(4).

In the degenerated case situation is analogous, only the Fermi energy plaices a part of the thermal energy, that is the equation for the point of stop is $U(r)\sim\varepsilon_F$. Let's use the method



FIG. 4: The potential energy of a gravitating particle in a cluster - the curves **(1)** and **(1')** and a particle in a polaron - the curves **(2)** and **(2')** at temperatures T_1 and T_2 accordingly. In the Boltzmann limit the radii of the cluster D and the polaron D_p is determined as the points of stop of a particle with the average kinetic energy kT, which explains these equality.

described in subsection (III C) in order to find radius of the polaron. This size is determined by the multiplier $\frac{2r_q}{\lambda^3}$ in Eq.(46), but we must get rid of temperature in the final expression. For this let's multiply $2r_q/\lambda^3$ to the dimensionless value kT/ε_F, where ε_F is Fermi energy (A7) determined by the average density of the system N/V. Then the polaron's radius is wrote as

$$\frac{1}{D_p^2} \propto \frac{r_q}{\lambda^3}\left(\frac{kT}{\varepsilon_F}\right) \propto \frac{q^2 m}{\hbar^2}\left(\frac{N}{V}\right)^{1/3}. \qquad (55)$$

This expression coincides with the expression for radius of cluster (44) without the correction for temperature in the system of attracting particles. Such coincidence is explained by the same as in classical case, but a part of the average kinetic energy kT plays Fermi energy, which we can understand as above-mentioned "statistical potential" too.

B. Hydrodynamical approach

In order to explain the coincidence of the radii of cluster D and polaron D_p we can use hydrodynamical approach [22] which lets to look at some general properties of system with Coulomb type interaction under other point of view. Let's consider the system of particle which is described be the persistent equation, Euler equation and Poisson equation ("+" for gravitating particles and "−" for interaction like electrical charges):

$$\begin{aligned} &\frac{\partial \rho}{\partial t} + \nabla \cdot \left(\rho \overrightarrow{v}\right) \\ \rho \frac{\partial \overrightarrow{v}}{\partial t} + \rho \left(\overrightarrow{v} \cdot \nabla\right) \overrightarrow{v} &= -\nabla P + \rho \nabla \phi \\ \Delta \phi &= \pm 4\pi G \rho. \end{aligned} \tag{56}$$

We suppose that perturbations is adiabatical: $P_1 = \frac{\partial P}{\partial \rho}\rho_1 = c^2 \rho_1$, where $c = \sqrt{kT/m}$ - the speed of sound, $P_1 \ll P$ - the small change of pressure. Then the linearized equation of motion for fluctuation ρ is

$$\left(\Delta - \frac{1}{c^2}\frac{\partial^2}{\partial t^2}\right)\rho_1 \pm k_J^2 \rho_1 = 0, \tag{57}$$

where $k_J = \frac{\sqrt{4\pi G \rho_0}}{c}$ - Jeans wave number.

Let's described localized oscillation process, when $k = 0$. Then we have next solution:

- For the system with Coulomb repulsion the dynamics of increase of the deflection has view $\rho_1 \propto \exp\left(i\omega t\right)$, where $\omega = ck = \sqrt{\frac{4\pi N m}{V}}$. That is we have plasma oscillations.
- for self-gravitation system the dynamics of increase of the deflection has view $\rho_1 \propto \exp\left(\omega t\right)$ or $\rho_1 \propto \exp\left(-\omega t\right)$, where $\omega = ck = \sqrt{\frac{4\pi N m}{V}}$.That means exponential increase of the deflection or relaxation to the equilibrium state - cluster.

We can see, that the both process is characterized by one and the same time scale $\tau \sim \frac{1}{\omega}/$ Then we can suppose, that the correspond spatial scale is distant, which a particle cross in the process of its thermal motion at the time$1/\omega$:

$$D \propto \frac{< v >}{\omega} = \sqrt{\frac{kT}{m}}/\omega = \sqrt{\frac{kTV}{4\pi G m^2 N}}. \tag{58}$$

This expression coincides with radiuses D and D_p for gravitating particles and repulsing particles accordingly. That is both aforesaid systems is characterized by one and the same sizes of spatial inhomogeneity.

As it has been said above, the partition function of self-gravitational system divergences, however the states exist where the equilibrium spatial distribution function is solution of Eq.(13). This means that our system disintegrates (collapses) to infinity multitude of clusters of size D_0 each, since the number of particles in the system $N \to \infty$, but the number of the particles in the cluster is terminal. This multitude is self-gravitating system too and it may be collapse. Then the process repeats again. In others words, the free anergy of the such system hasn't absolute minimum and each state of the self-gravitating system is analogous to false vacuum in theory of field [12]. To evaluate of times of relaxation we can using Eq.(58). So with each act of collapse, we must substitute $m \to nm$ and $N \to N/n$, where n is average number of particles in a cluster. Then the time scale $1/\omega$ is constant at each process of cluster formation (collapse).

V. CONCLUSION.

In this article, based on statistical approach [1, 2, 3, 4], we demonstrated a nonperturbative calculation of the partition function, we solved the system of particles with Boltzmann and Fermi statistics with long-range interaction of Coulomb type. All results have been obtained at the condition that average energy of interaction of two particles is much less than their average kinetic energy kT or Fermi energy ε_F: $\frac{q^2}{\sqrt[3]{V/N}} \ll kT$ or $\frac{q^2}{\sqrt[3]{V/N}} \ll \varepsilon_F$ - the "local ideality" approach [24].

The long-range attraction of particles results to the formation of the cluster of finite size, as the initial homogeneous state is instable in the limit $N \to \infty$, $V \to \infty$ with fixed N/V. The equation for spatial distribution function has be obtained. However, it is simplified in the next limit cases only .

In case of large temperature (when $\lambda \ll \sqrt[3]{v}$) the radius of cluster is determined by balance of the two energies: the energy of attraction interaction which aspires to compress gas, and the thermal energy which creates a positive pressure. This mechanism created equilibrium distribution in the the system (16), in particular, with rising of temperature the cluster's size enlarges and its central density decreases accordingly. In the case of degenerated system the radius of the cluster is determined by balance of the two energy too: energy of interaction which aspires to compress gas, and repulsive Fermi statistical potential creating positive pressure. In the total case the equilibrium cluster's size determined by three energy: energy of interaction, thermal and repulsive "statistical potential". The size of the cluster approaches to the equilibrium size asymptotically in process of its formation. The state of the system with spatially inhomogeneous distribution corresponding to the cluster of the equilibrium size (43 or 44) is stable if we are assuming that "local ideality" approach is true.

The size of spatial inhomogeneity of two matter: gravitating gas and repulsing particles (by Coulomb) is coincides at one and the same interaction's constant in Boltzmann limit and degenerated case. This coincidence caused by that, that the spatial distributions in both cases is determined by the two energies only: Coulomb type and thermal energy or repulsive Fermi "statistical potential". The mechanism of formation of the cluster and the polaron is one and the same - a balance of they two energies. This fact can be explained by hydrodynamical approach - both matters have same time scale of structure's formation, but in the first matter it determines time of relaxation and in the second matter it determines of plasma oscillation. This time scales determine spatial scale (the radii of the cluster and the polaron) monotonously.

APPENDIX A: THE EFFECTIVE FREE ENERGIES FOR IDEAL QUANTUM GASES.

Let's obtain some expressions used in this paper for ideal Fermi and Bose gas. Thus we'll demonstrate the correctness and greater rationality of our approach as compared with the traditional method [19].

In the case of ideal gas $\varphi = \psi = 0$. Then the effective free energy (3) for the system is

$$\begin{aligned} S = \quad & -\frac{1}{\omega}\int dV \int \; 4\pi p^2 dp \ln\left(1 + \xi \exp\left(-\beta\varepsilon_p\right)\right) + N\ln\xi \\ & = -\frac{V}{\lambda^3} f_{5/2}\left(\xi\right) + N\ln\xi, \end{aligned} \tag{A1}$$

where

$$\begin{aligned} f_{5/2}\left(\xi\right) \quad & = \frac{4}{\sqrt{\pi}}\int_0^\infty dx x^2 \ln\left(1 + \xi e^{-x^2}\right) \\ & = \sum_{l=1}^{\infty} (-1)^{l+1} \frac{\xi^l}{l^{5/2}} \end{aligned} \tag{A2}$$

is special Fermi function [19, 20]; $\varepsilon_p = p^2/2m$ is kinetic energy of a particle; $\lambda = \sqrt{\frac{\beta\hbar^2}{2\pi m}}$ is wave's thermal length of a particle. Then the equation for saddle point is as follows:

$$\frac{1}{v} = \frac{1}{\lambda^3} f_{3/2}\left(\xi\right), \tag{A3}$$

where $f_{3/2}\left(\xi\right) = \xi \frac{\partial f_{5/2}(\xi)}{\partial \xi}$, $v \equiv \frac{V}{N}$.

The partition function (2) for this case:

$$Z_N = \exp\left[\frac{V}{\lambda^3} f_{5/2}\left(\xi\right) - N\ln\xi\right], \tag{A4}$$

where ξ is determined from the equation (A3). We can find any thermodynamical functions knowing the partition function. Let's consider the case $\xi \to 0$ corresponding to high temperature (the Boltzmann limit), then Eqs. (A3,A4) are reduced to

$$Z_N = \exp\left[\frac{V}{\lambda^3}\xi - N\ln\xi\right] \approx \frac{V^N}{N!}\left(\frac{mkT}{2\pi\hbar^2}\right)^{\frac{3}{2}N} \tag{A5}$$

$$\frac{1}{v} = \frac{\xi}{\lambda^3} \tag{A6}$$

Let's consider the case $T \to 0$ corresponding to degenerated Fermi gas. Then the special Fermi functions are [19]

$$\begin{aligned} f_{3/2} &= \frac{4}{3\sqrt{\pi}}\left[(\ln\xi)^{3/2} + \frac{\pi^2}{8}(\ln\xi)^{-1/2}\right] + O(\xi^{-1}) \\ f_{5/2} &= \frac{4}{3\sqrt{\pi}}\left[\frac{2}{5}(\ln\xi)^{5/2} + \frac{\pi^2}{4}(\ln\xi)^{1/2}\right] + O(\xi^{-1}). \end{aligned} \tag{A7}$$

In this case, the expression (A3) is reduced to

$$\frac{\lambda^3}{v} \approx \frac{4}{3\sqrt{\pi}}(\ln\xi)^{3/2} \Rightarrow \xi \approx e^{\beta\varepsilon_F}, \tag{A8}$$

where

$$\varepsilon_F \equiv \frac{\hbar^2}{2m}\left(\frac{6\pi^2}{v}\right)^{\frac{2}{3}} \tag{A9}$$

is Fermi energy.

We can find analogical expressions for Bose gas. It is necessary to proceed from the effective free energy as

$$\begin{aligned} S = {} & \frac{1}{\omega}\int dV \int 4\pi p^2 dp \ln\left(1 - \xi\exp\left(-\beta\varepsilon_p\right)\right) + N\ln\xi \\ & = -\frac{V}{\lambda^3} g_{5/2}\left(\xi\right) + \ln\left(1-\xi\right) + N\ln\xi, \end{aligned} \tag{A10}$$

where $g_{5/2}\left(\xi\right) = -\frac{4}{\sqrt{\pi}}\int_0^\infty dx x^2 \ln\left(1 - \xi e^{-x^2}\right) = \sum_{l=1}^\infty \frac{\xi^l}{l^{5/2}}$ - special Bose function [19, 20]. It is clear that activity always $\xi < 1$ unlike Fermi system; $\ln\left(1-\xi\right)$ is effective free energy for condensed phase (the addendum with $\mathbf{p} = 0$ is as important as the rest of the sum when $\xi \to 1$).

Let's find the internal energy U of the system proceeding from the arranged analogy between apparatus of thermodynamics in our representation and the theory of field. In order to do it let's use the correlation $-H = \partial S_{mech}/\partial t$ and determine conformities $H \longleftrightarrow U, S_{mech} \longleftrightarrow S_{term}, t \longleftrightarrow 1/kT$, where H is Hamilton's function of the system, S_{mech} and S_{term} are the action for mechanics and effective free energy for thermodynamics (3) systems correspondingly, t is time and $1/kT$ is reverse temperature. Then, with the help of the expression (8), we have

$$U = -\frac{\partial S}{\partial\left(1/kT\right)} = \frac{3}{2}\frac{VkT}{\lambda^3} f_{5/2}\left(\xi\right) = \frac{3}{2}PV \tag{A11}$$

Thus the expressions for thermodynamical function of Fermi and Bose gases obtained by our method coincide with ones obtained by usual way [19, 20], and it confirms the correctness of the proposed approach.

REFERENCES

---

[1] E.D. Bilotsky, B.I. Lev, Teor. Math. Fiz. **60**, 120 (1984).

[2] E.D. Bilotsky, Doctor Degree Thesis, Institute of Physics, Kiev, 1992.

[3] B.I. Lev, A.Ya.Zhugaevych, Phys. Rev. E **57**, 6460 (1998).

[4] B.I. Lev, Phys. Rev. E **58**, R2681 (1998).

[5] V. Krasnoholovets, B. Lev, Cond. Matt. Phys **33** 1 (2002).

[6] K.V. Grigorishin and B.I. Lev, Phys. Rev. E **71**, 066105 (2005).

[7] B.I. Lev and K.V. Grigorishin, Cond. Matt. Phys **45** 67 (2006).

[8] J. Hubbard, Phys. Rev. Lett. **3**, 77 (1959).

[9] R.L. Stratonovich, Sov. Phys. Dokl. **2**, 416 (1984).

[10] S. Doniach, Phys. Rev. B **24**, 5063 (1981).

[11] P. Kampf, T.Zimanyi, Phys. Rev. B **47**, 279 (1993-I).

[12] S. Coleman, Phys. Rev. D **15**, 2929 (1977).

[13] R. Rajaraman, Phys. Rev. D **15**, 2866 (1977).

[14] L.N. Lipatov, Zn. Eksp. Teor. Fiz. **72**, 412 (1977).

[15] S. Edward and A.Lenard, J. Math. Phys. **3**, 778 (1962).

[16] S. Samuel, Phys. Rev. D **18**, 1916 (1978).

[17] V.B. Magalinsky, Zn. Eksp. Teor. Fiz. **48**, 167 (1965).

[18] R. Baxter, *Exactly Solved Models in Statistical Mehanics* (Academics Press, New York, 1982).

[19] K. Huang, *Statistical Mehanics* (Wiley & Sons, New York, 1963).

[20] A. Isihara, *Statistical Mehanics* (Academic, New York, 1971).

[21] L.D. Landau, E.M. Lifshits, *Quantum Mehanics* (Nauka, Moscow, 1989).

[22] J. Jeans, *Astronomy and Cosmogony* (Cambridge: Cambridge Univ. Press, 1928).

[23] H.J.Vega,N.Sanches and F.Combes,Phys.Rev.D.**54**, 6008,(1996)

[24] H.J.Vega, N.Sanches ,arXiv:hep-th/9903236 v2 (2000)

[25] I. Ispolatov and E.G.D. Cohen, Phys. Rev. Lett. **87**, 210601 (2001).

[26] M. Antoni and A. Torcini, Phys. Rev. E **57**, R6233 (1998).

[27] D. Ruelle, *Statistical Mehanics. Rigorous Results* (New York, Amsterdam, 1969).